

Where there is smoke: Solid fuel externalities, gender, and adult respiratory health in India

Aashish Gupta*

March 2, 2018

Abstract

Chronic respiratory conditions are the fourth biggest cause of death in the world. Using data on lung obstruction from the WHO Survey of Global AGEing and Adult Health (WHO-SAGE 2007-08), this paper studies the determinants of respiratory health in India, home to one-third of all deaths from Chronic Obstructive Pulmonary Disease. First, we find that smokers and members of households that use solid fuels (wood, biomass, coal or dung) for cooking have higher lung obstruction. Second, even if a respondent's household uses clean fuels, their lung obstruction is higher if their neighbours use solid fuels. In neighbourhoods with high solid fuel use, the lungs of members of households that use clean fuels can be as obstructed as lungs of members of households that use solid fuels. These negative externalities of solid fuel use are robust to additional controls for neighbourhood socioeconomic status, falsification tests, tests with placebo measures, and tests using alternative measures of respiratory health as outcomes, suggesting a causal link. Third, the influence of the determinants is patterned by gender. Smoking tobacco is an important influence on lung obstruction among men, while women from households that use solid fuels are the only group not further harmed by neighbourhood solid fuel smoke. The study improves our understanding of behavioural, social, and environmental determinants of respiratory health in India. Importantly, it makes a case for greater public investments to promote the adoption and use of cleaner fuels. (239 words)

Keywords: respiratory health, lung obstruction, solid fuels, neighbourhood context, negative externalities, gender, India

The paper includes a supplementary appendix for online publication. Word count, excluding the supplementary appendix, is 7485 words.

*Corresponding author. Email: aashishg@sas.upenn.edu. Population Studies Center & Department of Sociology, University of Pennsylvania. I am grateful to Jere Berhman, Jean Drèze, Michel Guillot, Emilio Parrado, Samuel Preston, Megan Reed, Alejandro Sánchez-Becerra, Dean Spears, Nikkil Sudharsanan, Harsha Thirumurthy, Atheendar Venkataramani, Yana Vierboom and especially to Diane Coffey, Irma Elo, Annette Lareau, and Kanika Sharma for helpful advice and comments. All errors are my own.

Where there is smoke: Solid fuel externalities, gender, and adult respiratory health in India

1 Introduction

Chronic respiratory conditions are the 4th largest cause of death in the world in 2016, responsible for 6.5 % of all deaths and 3.9 % of global Disability Adjusted Life Years lost ([GBD Causes of Death Collaborators, 2017](#)). Chronic Obstructive Pulmonary Disease (COPD), “a lung ailment that is characterized by a persistent blockage of airflow from the lungs” ([World Health Organisation, 2015](#)) accounts for 83% of all deaths from chronic respiratory conditions, and 68% of all Disability Adjusted Life Years lost. Although less developed countries have a higher burden of disease attributable to chronic respiratory diseases and adult respiratory health is becoming increasingly important for overall population health, empirical research on the determinants of respiratory health at the population level in developing countries is limited ([Mannino and Buist, 2007](#)). In high-income countries, smoking tobacco is the principle risk factor for COPD ([Currie, 2010](#); [Pauwels and Rabe, 2004](#)). Emerging evidence suggests that smoke from the use of solid fuels (wood, animal dung, coal, charcoal, and crop residues), which are used for cooking and heating, may be an equally important determinant in less developed countries ([Salvi and Barnes, 2009, 2010](#)).

Solid fuel smoke can harm the respiratory health of women ([Po et al., 2011](#); [Ezzati and Kammen, 2002](#)), as they are the household members primarily responsible for cooking ([Hirway and Jose, 2011](#)). Other household members can be harmed because of exposure to indoor air pollution ([Venkataramani and Fried, 2011](#); [Hu et al., 2010](#); [Larson and Rosen, 2002](#)). Researchers have hypothesised that because solid fuel smoke contributes to local air pollution, neighbours can be harmed as well, even if they themselves use clean fuels ([Smith, 2002](#); [Torres-Duque et al., 2008](#); [Balakrishnan et al., 2014](#); [Chafe et al., 2014](#)). However, existing research has not yet tested or quantified the association between neighbourhood externalities of solid fuel use and lung obstruction empirically. Additionally, while it is well-recognised that social, cultural and physical environments jointly influence health ([Elliott, 2017](#)), the patterning of the exposures by key social determinants such as gender has not been studied at the population level for developing countries.

This paper intends to fill these research gaps, for the important case of India.

Both smoking and cooking are patterned by gender in India, a large section of the Indian population is exposed to solid fuel smoke, and respiratory health is poor. Few women in India smoke ([Waldron et al., 1988](#); [Mishra et al., 2016](#)), while men do little household work or cooking ([Hirway and Jose, 2011](#)). Using data from the only representative survey that measured time use in India, [Hirway and Jose \(2011\)](#) found that while men spent less than an hour per week in India on cooking and related household activities, women spent an average of 25 hours per week. Thus, women who live in households that use solid fuels have high exposure to direct smoke while cooking ([Parikh et al., 1999](#)). More than 67.2% of Indian households used solid fuels for cooking ([Government of India, 2011a](#)), and about one-third of the world's population which relies on solid fuels was estimated to live in India ([Bonjour et al., 2013](#)). Finally, in 2016, India was estimated to be home to one-third of all deaths from chronic respiratory conditions ([GBD Causes of Death Collaborators, 2017](#)), and almost half of Indian adults had COPD ([World Health Organisation, 2013](#)) by the [GOLD \(2016\)](#) criteria. These factors make India is an important and urgent context to examine lung obstruction determinants.

We use population level survey data on lung obstruction collected by the World Health Organisation's Survey of Global AGEing and adult health (WHO-SAGE) 2007-08 ([Kowal et al., 2012](#)). The representative sample in the WHO-SAGE allows us to test for neighbourhood externalities, study the patterning of the risk-factors by gender, and examine the robustness of the results. From a social science perspective, this study contributes evidence from a developing country context to the literature on geography and health, as well as the literature on how social and behavioural determinants mediate this relationship ([Gatrell and Elliott, 2014](#); [Kawachi and Berkman, 2003](#)). From a health policy perspective, the study presents evidence of negative externalities, which are a classic argument for public intervention. We discuss policy and research implications of this study in section 5. The next section provides background information on lung obstruction. Section 3 discusses the measures and the empirical strategy, and section 4 presents results.

2 Background and significance

The lung is the body's organ of gas exchange. The trachea or the windpipe branches into the bronchi inside the lungs, which further divide into smaller branches, called bronchioles. The bronchioles end in microscopic air sacs called alveoli. Long-term exposure to noxious particles and gases produces inflammation in the lungs (Currie, 2010). This inflammation results in the narrowing of the bronchioles, and is accompanied by lung tissue destruction (emphysema), disruption of repair and defence mechanism in the lungs (bronchiolitis), and secretion of mucous and increased cough (chronic bronchitis) (Calverley and Walker, 2003). These changes increase resistance to airflow in the lungs. Persistent respiratory symptoms and lung obstruction characterise Chronic Obstructive Pulmonary Disease (GOLD, 2016). Lung obstruction can result in respiratory or heart failure (Holguin et al., 2005), increases the risk of lung infections and pneumonia (Prescott et al., 1995) while exacerbating the risk of death from co-morbidities such as heart disease and lung cancer (Sin et al., 2006).

Smoking tobacco is a well-established cause of lung obstruction. In a systematic review, Forey et al. (2011) pool odds-ratios from 133 studies and find higher odds of COPD in ever smokers, current smokers, and ex-smokers when compared to non-smokers. While smoking cessation cannot reverse damage, a systematic review found that compared to smoking, quitting was associated with slower increases in lung obstruction (Godtfredsen et al., 2008). Inhaling smoke from solid fuels is the other big risk factor for lung obstruction. Torres-Duque et al. (2008) and Kurmi et al. (2010) review evidence on the impact of solid fuel use on respiratory diseases, including COPD. Kurmi et al. (2010) find higher odds of having COPD among households that use solid fuels. In India, McKay et al. (2012); Moschovis et al. (2015); Dave et al. (2017) have found sex, smoking status, and indoor air pollution to be significant predictors of lung obstruction. However, some studies (Amaral et al., 2017; Sana et al., 2018) have failed to find significantly higher odds of COPD among those who used solid fuels. These studies have not considered solid fuel externalities. The comparison groups in these studies, those not using solid fuels, may also have high lung obstruction because of exposure to neighbourhood air pollution.

COPD is diagnosed using a non-invasive physiological test, spirometry, which measures lung obstruction. A spirometer measures the proportion of the volume of air (litres) one can exhale in

1 second to the total volume of air one exhales, after inhaling as deeply as possible (Cotes et al., 2009). This ratio is called Forced Expiratory Volume in 1 second percent (FEV1%). Higher values of FEV1% signify better respiratory health, and imply lesser obstruction of airflow. Existing research on solid fuel use and respiratory health relies on dichotomised COPD as the outcome, based on the Global initiative for Obstructive Lung Disease (GOLD) criteria of diagnosing COPD if FEV1% is less than 70% (GOLD, 2016). The GOLD cut-off over-diagnoses older patients, and under-diagnose younger patients (Pellegrino et al., 2005). While categorisation has clinical relevance, such dichotomisation does not capture the severity of the disease and reduces statistical power and precision (Weinberg, 1995; Royston et al., 2006). Consequently, we use continuous FEV1% as the outcome, but confirm that the results are robust to using dichotomised COPD.

The paper informs this literature by considering, for the first time, neighbourhood externalities of solid fuel use in an empirical framework, by using population representative survey data on lung obstruction, by examining the gendered patterning of the exposures, and by investigating both dichotomous and continuous measures of lung obstruction as outcomes. We do it for the important case of India, home to an estimated third of the total deaths from chronic respiratory conditions and COPD, and where COPD was also among the top three causes of death (Krishnan et al., 2011; Patel et al., 2011).

3 Materials and methods

3.1 Data and measures

The WHO-SAGE (2007-08) is representative of the adult population aged above 18 years in six states of India: Rajasthan, Uttar Pradesh, Assam, Karnataka, Maharashtra and West Bengal. These states had a combined population of 570 million in 2011, representing 47% of India's total population (Government of India, 2011b). The WHO SAGE conducted spirometry to measure lung obstruction for its 9,551 respondents (World Health Organisation, 2013), and collected information on other individual and household characteristics. Table 1 provides summary statistics for the main measures used in the analysis.

3.1.1 Outcome

The primary outcome is Forced Expiratory Volume in 1 second percent (FEV1%), the proportion of total air inhaled that a person can exhale in one second. FEV1% is a direct measurement of lung obstruction. The outcome is also called FEV1/FVC (ratio of Forced Expiratory Volume in 1 second to Forced Vital Capacity). Within clinical settings, FEV1% is often dichotomized, but the clinical criteria for diagnosis of COPD are contested ([Almagro et al., 2014](#); [Burney, 2014](#)). We use the continuous measure to preserve the full range of variation in lung function and improve the precision of our estimates. Our results were qualitatively unchanged when using a dichotomous outcome. Average FEV1% was 70.8% in the WHO-SAGE sample. Mean levels of FEV1% as low as these, very close to the GOLD criteria of FEV1% of 70% to clinically diagnose someone as having COPD, reinforce the impaired state of respiratory health in India. Men and women had similar mean values of lung obstruction (Table 1).

3.1.2 Exposures

The main independent variables that measure exposure to risk factors for lung obstruction are smoking; household solid fuel use, and neighbourhood solid fuel use.

Smoking: Smoking status was self-reported. Smoking is coded as a dummy variable (a very small fraction of the sample consisted of former smokers, and second-hand smoke was not observed). About 17% of the SAGE respondents smoked (Table 1). While 31% of the men smoked, only 2% of the women smoked.

Household use of solid fuels: Respondents were asked the fuel their household uses for cooking. Coal/charcoal, wood, agricultural residue/shrubs and animal dung were classified as solid fuels. Kerosene, gas, and electricity were classified as clean fuels. Household use of solid fuels is coded as a dummy variable. About 76% of all households used solid fuels for cooking.

Neighbourhood solid fuel use: Neighbourhood solid fuel use is measured as the proportion of households, excluding the respondent's household, in a Primary Sampling Unit (PSU) that use solid fuels. A PSU is an urban block or a rural village. Neighbourhood solid fuel use is coded as a continuous variable between 0 and 1. The SAGE survey interviewed respondents in 375 different clusters or PSUs, selecting on average 26 households per cluster. Although a larger sample from

each PSU would be desirable, and the neighbourhood solid fuel use measure may introduce measurement error due to extreme values (everyone using solid fuels or nobody using solid fuels), such specifications have been used productively in research on neighbourhood disease exposure, such as sanitation ([Geruso and Spears, forthcoming](#)). Solid fuel use in the neighbourhood is hypothesised to capture the deterioration in air quality in the neighbourhood due to solid fuel use. On average, 76% of a respondent's neighbours used solid fuels. Such a respondent would be assigned 0.76 as the value for the neighbourhood solid fuel use measure.

3.1.3 Other explanatory variables

In all regressions, we adjust for age dummies (in years), sex of the respondent, state fixed effects, and a dummy for living in a rural area. This specification of age has the advantage of accounting for non-linear relationships between age and lung obstruction. In the SAGE sample, average age was 40.4 years. Men were slightly older than women, and 47% of the sample was female. Almost 70% of the respondents lived in rural areas (Table 1). Apart from demographic and place controls, we also control for two measures of socioeconomic status, namely, years of education and number of household assets owned. We use dummies for years of education and count of assets, a stronger control than linear controls for wealth and education. Years of education ranged from 0 to 18. Mean years of education were 5.9 years, 7.4 years for men and 4.2 years for women, reflecting an aspect of gender inequality well-known for India ([Drèze and Sen, 2013](#)). The WHO SAGE asked respondents the number of chairs, tables, and cars their household owned, whether the household had a cycle, clock, bucket, cot, fridge, fixed line phone, mobile phone, television, computer, radio, livestock, sewing machine, a two-wheeler, bullock cart, and non-homestead agricultural land. Based on these indicators, we created a variable for the count of assets a respondent's household owned. The mean number of assets was 13.2.

3.2 Empirical strategy

The empirical strategy consists of several steps, including descriptive graphs of cumulative density function of lung obstruction by exposure type, Ordinary Least Squares (OLS) regressions, and

multiple robustness checks. The OLS linear regression equation is specified as:

$$\begin{aligned} \text{respiratory health}_{ij} = & \beta_1 \text{smoke tobacco}_{ij} + \beta_2 \text{household solid fuel use}_{ij} + \beta_3 \text{neighbourhood solid fuel use}_{ij} \\ & + \alpha \text{place}_{ij} + \theta \text{demographic}_{ij} + \lambda \text{ses}_{ij} + \varepsilon_{ij}, \end{aligned} \tag{1}$$

where respiratory health is a measure of obstruction of airflow of respondent i living in primary sampling unit j , measured as FEV1%. We cluster standard errors at the level of the primary sampling unit, to account for the WHO SAGE's multi-stage sample design, and use survey weights throughout. The coefficients of interest are β_1 , of whether the respondent smokes, β_2 , of whether the respondent's household uses solid fuels, and β_3 , of the proportion of other households in a respondent's primary sampling unit who use solid fuels. β_3 is a continuous measure between 0 and 1. Since regression coefficients show the effect of a one-unit change, β_3 indicates the difference between respondents who live in neighbourhoods where no other households use solid fuels and respondents who live in neighbourhoods where all other households use solid fuels.

place_{ij} controls for state fixed effects and a dummy for rural residence; demographic_{ij} adjusts for sex of the respondent and age dummies, in years; and ses_{ij} controls include dummies for both years of education and count of household assets.

In addition to these controls for SES variables, we conduct further robustness checks, adjusting for additional household and neighbourhood SES variables, such as log household expenditure and average neighbourhood assets. In placebo tests, we test whether measures similar to the exposures, such as chewing tobacco, drinking alcohol, household ownership of toilets, household electrification, neighbourhood open defecation or neighbourhood electrification are associated with the outcome. In falsification tests, we test whether other health outcomes, such as Forced Vital Capacity of lungs or adult heights are associated with the exposures. Finally, we test if the exposures are associated with alternative measures of respiratory health, such as self-reported lung disease and Forced Expiratory Volume in 1 second. In placebo tests, it is expected that there will be no association between the placebo variables and the outcome. In falsification tests, it is expected that there will be no association between the falsification outcomes and the exposures.

We expect to find an association between the alternative measures of respiratory health and the exposures. These results are available in the supplementary appendix.

4 Results

The results section is organised as follows. We begin by descriptive graphs in section 4.1, first examining the cumulative density function of FEV1% by the three exposures. We then see the extent to which the differences in the exposures can be explained by socioeconomic status. Thereafter, we examine whether household and neighbourhood solid fuel use jointly influence lung obstruction. Section 4.2 presents regression results, documenting the association between the outcome and the exposures while adjusting for other explanatory variables, and examining the implications of the gendered patterning of the exposures. Finally, section 4.4 presents the results from robustness checks, including the placebo and falsification tests, and tests with alternative measures of respiratory health as outcomes.

4.1 Descriptive graphs

Figure 1 presents cumulative density function of FEV1% for those exposed to the three risk-factors of lung obstruction (smoking, household solid fuel use, neighbourhood solid fuel use) and those not exposed. The cumulative distributions show differences in lung obstruction by exposure at all levels of the distribution of FEV1%. p -values and D -values (absolute maximum distance between the two distributions) from Kolmogorov-Smirnov tests, and p -values and z -statistics from Wilcoxon rank-sum test are also shown to compare the distributions in each of the sub-figures. The p values, which are all less than .001, indicate that the cumulative density functions of those exposed to the risk factors and those not exposed are significantly different. Smokers, members of households that use solid fuels, and residents of neighbourhoods in which other members use solid fuels have higher lung obstruction across the distribution, than non-smokers, members of households that use clean fuels and residents of neighbourhoods that have lower levels of solid fuel use, respectively. The figure reveals that differences by household and neighbourhood solid fuel use exposure are larger than or similar to the differences by smoking.

Figure 2 provides graphical evidence that these differences cannot be explained by differences

in socioeconomic status. In fact, measures of socioeconomic status, such as years of education and household assets themselves have a weak relationship with lung obstruction. In Figure 2 the left panel shows differences in years of education, and the right panel shows differences by count of household assets using non-parametric local polynomial regressions with 95% confidence intervals shaded blue. At all levels of education and household assets, respondents whose households or neighbours use solid fuels have higher lung obstruction and lower values of FEV1%.

Figure 3 shows differences in lung obstruction between members of households that use solid fuels and members of households that use clean fuels by neighbourhood solid fuel use. The x-axis shows the proportion of households in a primary sampling unit using solid fuels, and the y-axis shows FEV1%. Among those respondents whose households use clean fuels, there is a much steeper decline in FEV1% as neighbourhood solid fuel use rises. Among those respondents whose households use clean fuels, as neighbourhood solid fuel increases, a modest decline in FEV1% can be seen. That respondents who use cleaner fuels show a steeper decline in FEV1% as neighbourhood solid fuel use increases reinforces existing evidence of a downward sloping concave concentration response function, which implies diminishing marginal impacts of increases in pollution (Burnett et al., 2014; Pope III et al., 2015). As does the shape of the non-linear decline among respondents whose households use clean fuels. Respondents already exposed to direct or indoor smoke from solid fuel use are not much harmed by neighbourhood solid fuels.

4.2 Regression results

Table 2 provides results from OLS regressions. The results indicate that average differences by the exposures are statistically significant, in magnitude similar to the ones in the descriptive graphs, and robust to demographic, place, and SES controls. The first 3 models present results for the overall sample, models 4 and 5 restrict the sample to women, and models 6 and 7 consider only men. All regressions adjust for age dummies, sex, state fixed effects and rural residence. All models except model 1 also adjust for socioeconomic status, by using dummies for years of education and number of household assets owned. Models with linear controls for age and measures of socioeconomic status are available in appendix table A1. Appendix table A1 shows that assets and education are not associated with lung obstruction, and that dummies for age and socioeconomic

status improve model fit.

Model 1 of table 2 shows that those who smoke, whose household uses solid fuels, and whose neighbours use solid fuels have lower FEV1% and higher lung obstruction. Model 2 adjust for socioeconomic status but the point estimates or the significance of the coefficients do not change substantially. Compared to a respondent who does not smoke, whose household does not use solid fuels, and whose neighbours do not use solid fuels, a respondent who smokes, or a respondent whose household uses solid fuels, is, on average, able to breathe out 3.7 percentage points less oxygen in the first second. Holding smoking and household solid fuel use constant, a respondent living in a neighbourhood in which all his neighbours use solid fuels is able to breathe out 6 percentage points less FEV1%. Models 1 and 2 reinforce the results from the descriptive graphs, that household solid fuel use and smoking tobacco are equivalent risks, and neighbourhood solid fuel use is an even greater risk for lung obstruction. In other words, a person who is a member of a household that does not use solid fuels, does not smoke, but lives in a neighbourhood in which half the households use solid fuels is likely to have the same level of lung obstruction as a person who is a tobacco smoker.

Model 3 shows the interaction between household and neighbourhood solid fuel use. Predicted FEV1% for household and neighbourhood solid fuel use are available in figure 4. The interaction results and predicted FEV1% closely replicate the results from figure 3: as neighbourhood solid fuel rises, FEV1% declines more steeply among members of households that use solid fuels.

4.3 Gender, exposures, and lung obstruction

Women and men differ both in the extent of their exposure to solid fuel smoke and smoking tobacco (Waldron et al., 1988; Hirway and Jose, 2011), as well as their physiological responses to the exposures (Aryal et al., 2013), making a case for within-sex comparisons. The next set of models in table 2 present stratified regressions for women (models 4 and 5) and men (models 6 and 7).

Model 4 of table 2 shows that while smoking has a negative association with FEV1% among women, the effect size is much smaller than that of men (model 7), and not significant. This is

likely due to selection, lesser intensity of smoking among women, a very small number of women for whom smoking is observed, and possible reporting bias because women may be less willing to say that they smoke. Household solid fuel smoke is significantly and negatively associated with FEV1% among women. While neighbourhood solid fuel use has a negative association with FEV1%, this effect isn't statistically significant (p -value 0.128). However, model 5, which interacts household and neighbourhood solid fuel use for women, shows that neighbourhood solid fuel use has a significantly negative association for women whose households don't use solid fuels. On the other hand, women who are members of households that use solid fuels are not further harmed by neighbourhood solid fuel use. This result can be more clearly seen through predicted margins, reported in figure 4.

Among men, smoking is associated with higher lung obstruction and lower FEV1%. The association of household use of solid fuels has a similar magnitude as the association of smoking, while the neighbourhood solid fuel use coefficient is even higher than that of smoking (model 6). Model 7 reports coefficients from an interaction of household solid fuel use and neighbourhood solid fuel use. For men, the interaction is not significant. These patterns, consistent with [Burnett et al. \(2014\)](#) and [Pope III et al. \(2015\)](#), underscore the high levels of exposure to smoke among women exposed to solid fuel smoke from direct cooking.

4.4 Robustness tests

As described previously, we next examine the robustness of the results. The regression tables are shown in the supplementary appendix.

Table [A2](#) adds controls for log monthly expenditure, log monthly expenditure squared, average neighbourhood assets, and average neighbourhood assets sequentially to the main specification from Table 2 (model 2), repeated as Model 1 in appendix table [A2](#) for comparison. The log of monthly household expenditure and the square of the log expenditure are not significantly associated with lung obstruction (model 2). Nor does the inclusion of these controls change the magnitude of the associations for the three exposures. Models 3 and 4 add controls for average neighbourhood assets (model 3) and average neighbourhood education (model 4). As with neighbourhood solid fuel use, these variables are constructed as average of other households' average

assets, or other respondents' average education. They are also not associated with FEV1%, and do not substantively change the effect sizes of the exposures.

Table A3 reports placebo tests for smoking tobacco, household solid fuel use, and neighbourhood solid fuel use. The associations of chewing tobacco and drinking alcohol with FEV1% are compared to that of smoking tobacco. Owning a toilet and household electrification, measures of household SES, are used as placebos for solid fuel use. Measures for neighbourhood electrification and neighbourhood open defecation are used as placebos for neighbourhood solid fuel use. As before, all regressions control for demographic, place, and other socioeconomic status variables. Panel A shows that there is no association between FEV1% and chewing tobacco, or between FEV1% and drinking alcohol. Chewing tobacco and drinking alcohol cause other health problems, but are not linked to respiratory health, either theoretically or empirically in the WHO-SAGE. Regressions in Panel B find no association of FEV1% with household ownership of toilets and household electrification, and regressions in Panel C find no association of lung obstruction with neighbourhood electrification or neighbourhood open defecation. The effect sizes for the placebo variables are also close to zero.

In Table A4, we conduct falsification tests (models 1 and 2) and tests with alternative measures of respiratory health (models 3 - 5). It is expected that the exposures will not be associated with falsification outcomes, but will be associated with alternative measures of respiratory health. It is well known that exposures such as smoking or solid fuel use harm airflow obstruction but not total lung capacity, which we confirm in model 1. The outcome in model 2 is height of the respondents. While height is influenced by early childhood disease environment, adult exposures to solid fuel use should not influence height, which is confirmed in model 2. Model 3 shows that smoking tobacco, household use of solid fuels, and neighbourhood solid fuel use are negatively associated with Forced Expiratory Volume in 1 second, measured in litres. Because the amount of air blown out is also associated with stature, model 3 adjusts for height in meters. Model 4 and 5 use the logistic model with binary outcomes of having COPD (following GOLD 2016 guidelines) and self-reported symptoms of lung disease, following the WHO SAGE guidelines (Arokiasamy et al., 2015). Both models show higher odds of having COPD and self-reported lung diseases among smokers, members of households who use solid fuels, and residents of neighbourhoods

with high solid fuels use, reinforcing the main finding.

5 Discussion

This paper expands evidence on the determinants of lung obstruction for the important case of India, where one-third of all deaths due to chronic respiratory conditions take place, in three key ways. First, to our knowledge, this is the first study that documents an empirical association between local solid fuel use by other households in one's neighbourhood and obstruction in one's own lungs. Multiple robustness checks suggest a causal interpretation of this association. Second, this study improves our understanding of gender as a determinant of respiratory health in India. We find that women who live in households that use solid fuels are not further harmed by neighbourhood solid fuel use. They are the only group not further harmed by neighbourhood solid fuel use, which points to the already high exposure to indoor smoke they must have while cooking. Gender also influences smoking behaviour. Finally, the study establishes that large sections of the Indian population - those who smoke, those who use solid fuels, and those who live near others who use solid fuels - have higher lung obstruction than those who aren't exposed to these determinants of lung obstruction. Higher socioeconomic status does not mitigate the influence of these exposures.

From the point of view of estimating differences due to the exposures over a person's life course, the study offers conservative estimates. Some of the respondents who were not exposed at the time of the survey would have been exposed to the exposures earlier in their lives. However, it has some limitations. The lungs of women who cook with solid fuels are more likely to be obstructed ([Johnson et al., 2011](#)), but cooking behaviour is not observed in the WHO-SAGE. Future health surveys should identify household members responsible for cooking, and if possible, ask about duration of cooking, so that dose-response relationships can be understood. We are unable to include measures of pollution in our analysis, as pollution monitoring in India is in a nascent stage ([Guttikunda, 2017](#)).

The study is also limited because it uses cross-sectional data. However, from a policy-makers' perspective, this study elevates the importance of promoting clean fuels to improve adult health in India. Even though currently exposed populations may not see immediate or significant im-

improvements in FEV1% (Chapman et al., 2005; Smith-Sivertsen et al., 2009; Kurmi et al., 2012), further obstruction can be prevented. Encouraging the adoption of cleaner fuels remains a non-trivial and hard challenge (Mobarak et al., 2012; Peel et al., 2015; Jeuland et al., 2015). But the only viable public health strategy for reducing lung obstruction is prevention through the adoption of cleaner fuels. Existing government programmes in India to promote cleaner fuels are means-tested, provide modest capital subsidies, involve significant private investments, do not have components for encouraging behaviour change, and focus on private harms to women from cooking with solid fuels (Ministry of Petroleum and Natural Gas, 2016; Sethi and Deep, 2018). There is a case for greater investments in efforts to communicate the private and neighbourhood harms of solid fuel use and encouraging households to switch to cleaner fuels.

From a population health research perspective, there is an urgent need for research that integrates social determinants, environmental exposures, measures of pollution, and respiratory health outcomes. Existing evidence points that women's lower status impedes the adoption of cleaner fuels in India (Kishore and Spears, 2014). Researching the role of gender in the persistence of solid fuel use in India, despite fast economic growth, and investigating approaches that address this social determinant of adoption of cleaner fuels are an important research concern as well.

References

- Almagro, Pere et al.** 2014. "Finding the best thresholds of FEV1 and dyspnea to predict 5-year survival in COPD patients: the COCOMICS study." *PloS one*, 9(2): e89866.
- Amaral, André FS et al.** 2017. "Airflow obstruction and use of solid fuels for cooking or heating: BOLD results." *American journal of respiratory and critical care medicine*(ja).
- Arokiasamy, Perianayagam et al.** 2015. "The impact of multimorbidity on adult physical and mental health in low-and middle-income countries: what does the study on global ageing and adult health (SAGE) reveal?" *BMC medicine*, 13(1): 178.
- Aryal, Shambhu et al.** 2013. "COPD and gender differences: an update." *Translational Research*, 162(4): 208–218.
- Balakrishnan, Kalpana et al.** 2014. "Addressing the burden of disease attributable to air pollution in India: the need to integrate across household and ambient air pollution exposures." *Environmental health perspectives*, 122(1): A6.
- Bonjour, Sophie et al.** 2013. "Solid fuel use for household cooking: country and regional estimates for 1980-2010." *Environmental Health Perspectives (Online)*, 121(7): 784.
- Burnett, Richard T et al.** 2014. "An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure." *Environmental health perspectives*, 122(4): 397.
- Burney, Peter.** 2014. "Coming off the GOLD standard." *The Lancet Respiratory Medicine*, 2(3): 174–

176.

- Calverley, PMA and Paul Walker.** 2003. "Chronic obstructive pulmonary disease." *The Lancet*, 362(9389): 1053 – 1061.
- Chafe, Zoë A et al..** 2014. "Household cooking with solid fuels contributes to ambient PM_{2.5} air pollution and the burden of disease." *Environmental health perspectives*, 122(12): 1314.
- Chapman, Robert S et al..** 2005. "Improvement in household stoves and risk of chronic obstructive pulmonary disease in Xuanwei, China: retrospective cohort study." *Bmj*, 331(7524): 1050.
- Cotes, John E et al..** 2009. *Lung function: physiology, measurement and application in medicine*: John Wiley and Sons.
- Currie, Graeme P.** 2010. *ABC of COPD*: John Wiley & Sons.
- Dave, Mona et al..** 2017. "Household air pollution and lung function in Indian adults: a cross-sectional study." *The International Journal of Tuberculosis and Lung Disease*, 21(6): 702–704.
- Drèze, Jean and Amartya Sen.** 2013. *An uncertain glory: India and its contradictions*: Princeton University Press.
- Elliott, Susan J.** 2017. "Environment and Health." *International Encyclopedia of Geography: People, the Earth, Environment and Technology*.
- Ezzati, Majid and Daniel M Kammen.** 2002. "The health impacts of exposure to indoor air pollution from solid fuels in developing countries: knowledge, gaps, and data needs.." *Environmental health perspectives*, 110(11): 1057.
- Forey, Barbara A et al..** 2011. "Systematic review with meta-analysis of the epidemiological evidence relating smoking to COPD, chronic bronchitis and emphysema." *BMC pulmonary medicine*, 11(1): 1.
- Gatrell, Anthony C and Susan J Elliott.** 2014. *Geographies of health: An introduction*: John Wiley & Sons.
- GBD Causes of Death Collaborators.** 2017. "Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016." *Lancet*, 390(10100): 1151–1210.
- Geruso, Michael and Dean Spears.** forthcoming. "Neighborhood Sanitation and Infant Mortality." *American Economic Journal: Applied Economics*.
- Godtfredsen, NS et al..** 2008. "COPD-related morbidity and mortality after smoking cessation: status of the evidence." *European Respiratory Journal*, 32(4): 844–853.
- GOLD.** 2016. "Global Strategy for the Diagnosis, Management and Prevention of COPD." Technical report, Global Initiative for Chronic Obstructive Lung Disease - GOLD.
- Government of India.** 2011a. "Census of India 2011: Household Listing Operations." *Office of the Registrar General Census Commissioner, India*.
- 2011b. "Census of India 2011: Houselisting Operations." *Housing and Houselisting Operations*.
- Guttikunda, Sarath.** 2017. "Its About Time We Got Smarter About Monitoring Our Air Pollution." *The Wire*.
- Hirway, Indira and Sunny Jose.** 2011. "Understanding women's work using time-use statistics: The case of India." *Feminist Economics*, 17(4): 67–92.
- Holguin, Fernando et al..** 2005. "Comorbidity and mortality in COPD-related hospitalizations in the United States, 1979 to 2001." *Chest Journal*, 128(4).
- Hu, Guoping et al..** 2010. "Risk of COPD from exposure to biomass smoke: a metaanalysis." *Chest Journal*, 138(1): 20–31.
- Jeuland, Marc et al..** 2015. "The economics of household air pollution." *Annu. Rev. Resour. Econ.*, 7(1): 81–108.

- Johnson, Priscilla et al.**. 2011. "Prevalence of chronic obstructive pulmonary disease in rural women of Tamilnadu: implications for refining disease burden assessments attributable to household biomass combustion." *Global health action*, 4(1): 7226.
- Kawachi, Ichiro and Lisa F Berkman.** 2003. *Neighborhoods and health*: Oxford University Press.
- Kishore, Avinash and Dean Spears.** 2014. "Having a son promotes clean cooking fuel use in urban India: Womens status and son preference." *Economic Development and Cultural Change*, 62(4): 673–699.
- Kowal, Paul et al.**. 2012. "Data resource profile: the World Health Organization Study on global AGEing and adult health (SAGE)." *International journal of epidemiology*, 41(6): 1639–1649.
- Krishnan, Anand et al.**. 2011. "Adult mortality surveillance by routine health workers using a short verbal autopsy tool in rural north India." *Journal of Epidemiology & Community Health*.
- Kurmi, Om P et al.**. 2010. "COPD and chronic bronchitis risk of indoor air pollution from solid fuel: a systematic review and meta-analysis." *Thorax*, 65(3): 221–228.
- . 2012. "Indoor air pollution and the lung in low-and medium-income countries." *European Respiratory Journal*, 40(1): 239–254.
- Larson, Bruce A and Sydney Rosen.** 2002. "Understanding household demand for indoor air pollution control in developing countries." *Social science & medicine*, 55(4): 571–584.
- Mannino, David M and A Sonia Buist.** 2007. "Global burden of COPD: risk factors, prevalence, and future trends." *The Lancet*, 370(9589): 765–773.
- McKay, Ailsa J et al.**. 2012. "Prevalence of COPD in India: a systematic review." *Primary Care Respiratory Journal*, 21: 313–321.
- Ministry of Petroleum and Natural Gas.** 2016. "Pradhan Mantri Ujjwala Yojana - modalities for implementation." Technical report, Government of India.
- Mishra, Sujata et al.**. 2016. "Trends in bidi and cigarette smoking in India from 1998 to 2015, by age, gender and education." *BMJ global health*, 1(1).
- Mobarak, Mushfiq et al.**. 2012. "Low demand for nontraditional cookstove technologies." *Proceedings of the National Academy of Sciences*, 109(27): 10815–10820.
- Moschovis, Peter P et al.**. 2015. "The Association Of Household Air Pollution And Smoking With Risk Of Obstructive Lung Disease Among Adults In India." in *A105. COPD EPIDEMIOLOGY: GLOBAL PERSPECTIVE*: Am Thoracic Soc: A6353–A6353.
- Parikh, Jyoti et al.**. 1999. "Indoor air pollution: a reflection on gender bias." *Economic and Political Weekly*: 539–544.
- Patel, Vikram et al.**. 2011. "Chronic diseases and injuries in India." *The Lancet*, 377(9763): 413–428.
- Pauwels, Romain A and Klaus F Rabe.** 2004. "Burden and clinical features of chronic obstructive pulmonary disease (COPD)." *The Lancet*, 364(9434): 613–620.
- Peel, Jennifer L et al.**. 2015. "Are Randomized Trials Necessary to Advance Epidemiologic Research on Household Air Pollution?" *Current Epidemiology Reports*, 2(4): 263–270.
- Pellegrino, Riccardo et al.**. 2005. "Interpretative strategies for lung function tests." *European Respiratory Journal*, 26(5): 948–968.
- Po, June YT et al.**. 2011. "Respiratory disease associated with solid biomass fuel exposure in rural women and children: systematic review and meta-analysis." *Thorax*, 66(3): 232–239.
- Pope III, C Arden et al.**. 2015. "Health benefits of air pollution abatement policy: role of the shape of the concentration–response function." *Journal of the Air & Waste Management Association*, 65(5): 516–522.
- Prescott, E et al.**. 1995. "Chronic mucus hypersecretion in COPD and death from pulmonary infection." *European Respiratory Journal*, 8(8): 1333–1338.
- Royston, Patrick et al.**. 2006. "Dichotomizing continuous predictors in multiple regression: a bad

- idea." *Statistics in medicine*, 25(1): 127–141.
- Salvi, Sundeep S and Peter J Barnes.** 2009. "Chronic obstructive pulmonary disease in non-smokers." *The lancet*, 374(9691): 733–743.
- . 2010. "Is exposure to biomass smoke the biggest risk factor for COPD globally?" *CHEST Journal*, 138(1): 3–6.
- Sana, Adama et al..** 2018. "Chronic obstructive pulmonary disease associated with biomass fuel use in women: a systematic review and meta-analysis." *BMJ Open Respiratory Research*, 5(1): e000246.
- Sethi, Nitin and Aroon Deep.** 2018. "PMs plan for free gas connections is failing its objective as government had been warned it would." *Scroll.in*.
- Sin, Don D et al..** 2006. "Mortality in COPD: role of comorbidities." *European Respiratory Journal*, 28(6): 1245–1257.
- Smith, Kirk R.** 2002. "Indoor air pollution in developing countries: recommendations for research." *Indoor air*, 12(3): 198–207.
- Smith-Sivertsen, Tone et al..** 2009. "Effect of reducing indoor air pollution on women's respiratory symptoms and lung function: the RESPIRE Randomized Trial, Guatemala." *American journal of epidemiology*, 170(2): 211–220.
- Torres-Duque, Carlos et al..** 2008. "Biomass fuels and respiratory diseases: a review of the evidence." *Proceedings of the American Thoracic Society*, 5(5): 577–590.
- Venkataramani, Atheendar S and Brian J Fried.** 2011. "Effect of worldwide oil price fluctuations on biomass fuel use and child respiratory health: evidence from Guatemala." *American journal of public health*, 101(9): 1668–1674.
- Waldron, Ingrid et al..** 1988. "Gender differences in tobacco use in Africa, Asia, the Pacific, and Latin America." *Social science & medicine*, 27(11): 1269–1275.
- Weinberg, Clarice R.** 1995. "How bad is categorization?" *Epidemiology*: 345–347.
- World Health Organisation.** 2013. *Study on Global AGEing and Adult Health: India*: WHO & IIPS.
- . 2015. "World Health Organisation Fact-Sheet on Chronic Obstructive Pulmonary Disease." *Geneva: World Health Organization*.

Table 1: Descriptive statistics for the sample

	Total		Women		Men	
	mean	s.e.	mean	s.e.	mean	s.e.
Outcome						
Forced Expiratory Volume 1s %	70.8	0.5	71.2	0.7	70.5	0.6
Exposures						
Respondent smokes	0.17	0.01	0.02	0.00	0.31	0.02
Household uses solid fuels	0.76	0.02	0.77	0.02	0.76	0.02
Neighbourhood solid fuel use	0.76	0.02	0.77	0.02	0.76	0.02
Explanatory variables						
<i>Demographic</i>						
Age	40.4	0.3	39.4	0.3	41.3	0.4
Female	0.47					
<i>Socio-economic Status</i>						
Years of education	5.9	0.2	4.2	0.2	7.4	0.2
# of assets owned	13.2	0.2	13.2	0.2	13.3	0.2
<i>Contextual</i>						
Rural	0.70	0.02	0.70	0.03	0.70	0.02
N	9551		5777		3774	

Source: WHO SAGE 2007-08

Note: All means are weighted using national individual weights, and standard errors are clustered at the level of the primary sampling unit. FEV1% = (FEV1s * 100) / FVC.

Table 2: OLS regression coefficients of the association of smoking, household solid fuel use, and neighbourhood solid fuel use with lung obstruction




	overall sample			women		men	
	(1)	(2)	(3)	(4)	(5)	(6)	(7)
smoke tobacco	-3.403** (1.04)	-3.764*** (1.04)	-3.812*** (1.04)	-0.307 (3.183)	-0.506 (3.185)	-3.510** (1.130)	-3.529** (1.131)
household uses solid fuels	-3.226* (1.25)	-3.762** (1.17)	-7.407* (3.01)	-3.484* (1.460)	-10.510* (4.171)	-4.166* (1.725)	-5.460 (3.999)
neighbourhood solid fuel use	-6.082* (2.50)	-6.088* (2.45)	-7.299*** (2.06)	-3.399 (2.231)	-8.831*** (2.480)	-5.259* (2.457)	-6.153+ (3.453)
hh X neighbourhood solid fuel use			5.79 (3.71)		11.091* (5.149)		2.011 (5.191)
female	-0.396 (0.84)	-0.859 (0.91)	-0.849 (0.91)				
demographic controls	X	X	X	X	X	X	X
place controls	X	X	X	X	X	X	X
ses controls		X	X	X	X	X	X
constant	72.829*** (1.96)	74.768*** (2.62)	75.497*** (2.71)	71.536*** (3.284)	73.507*** (3.490)	76.119*** (3.151)	76.403*** (3.355)
observations	9551	9551	9551	5777	5777	3774	3774

Source: WHO SAGE 2007-08

Note: Robust standard errors, clustered at the level of the primary sampling unit, in parentheses. + p<.1, * p<.05, ** p<.01, *** p<.001.


Outcome: FEV1% (FEV1s/FVC as a percentage) is the ratio of the volume of air (litres) one can exhale in 1 second to the total volume of air one can exhale (after inhaling as deeply as possible). Demographic controls: age dummies and sex (coefficients for female shown). Place controls: rural residence and state fixed effects. SES controls: dummies for years of education and count of assets. Appendix Table 1 compares models with linear controls for age and ses measures with fixed effects for age and ses measures. Appendix Figure 1 shows the predicted margins of the interaction of household and neighbourhood solid fuel use, from models 3, 5, and 7. hh: household.

Figure 1: Cumulative Density Function of lung obstruction by exposure type




Source: SAGE 2007-08. Outcome: FEV1% (FEV1s/FVC as a percentage) is the ratio of the volume of air (liters) one can exhale in 1 second to the total volume of air one can exhale (after inhaling as deeply as possible). Higher values denote lesser lung obstruction. hh: household

Figure 2: Lung obstruction by household and neighbourhood exposure, and measures of Socio-economic Status




Source: SAGE 2007-08. Outcome: FEV1% (FEV1s/FVC as a percentage) is the ratio of the volume of air (liters) one can exhale in 1 second to the total volume of air one can exhale (after inhaling as deeply as possible). Higher values denote lesser lung obstruction. hh: household

Figure 3: Lung obstruction by neighbourhood solid fuel use among households that use solid fuels and households who use clean fuels



Source: SAGE 2007-08. Outcome: FEV1% (FEV1s/FVC as a percentage) is the ratio of the volume of air (liters) one can exhale in 1 second to the total volume of air one can exhale (after inhaling as deeply as possible). Higher values denote lesser lung obstruction. hh: household

Figure 4: Predicted FEV1% from OLS regressions with household and neighbourhood solid fuel interactions



Source: SAGE 2007-08. Outcome: FEV1% (FEV1s/FVC as a percentage) is the ratio of the volume of air (liters) one can exhale in 1 second to the total volume of air one can exhale (after inhaling as deeply as possible). Higher values denote lesser lung obstruction. Graphs show local polynomial regression. hh: household

SUPPLEMENTARY APPENDIX FOR ONLINE PUBLICATION

Appendix table [A1](#) examines the association of smoking, household solid fuel use, and neighbourhood solid fuel use with lung obstruction. The models in the table compare the association of the exposures when considered individually (models 1 - 3) and when considered together (models 4 - 6), the magnitudes of association for linear controls of age and socioeconomic status (models 1 - 5), and the magnitude of association for living in a rural area (models 1 - 6). Measures of socioeconomic status are not associated with lung obstruction. Model 6 in [A1](#) is the same as Model 2 in table [2](#). Operationalising the measures of socioeconomic status and age as dummies improves model fit, as shown by the r^2 . The table shows coefficients for the rural dummy as well, finding that respondents in rural areas are not significantly different from urban respondents once household and neighbourhood solid fuel use are adjusted for.

Appendix Tables [A2](#) - [A4](#) examine the robustness of the main result with controls for additional measures of socioeconomic status, placebo tests, falsification tests, and tests with alternative measures of respiratory health. The main findings from these tables are discussed in section [4.4](#).

Table A1: Association between smoking, household use of solid fuels, and neighbourhood solid fuel use with lung obstruction: Models with linear controls for age, assets, and education

	(1)	(2)	(3)	(4)	(5)	(6)
smoke tobacco	-3.909*** (1.069)			-3.484** (1.066)	-3.673*** (1.062)	-3.764*** (1.040)
household uses solid fuels		-5.241*** (1.427)		-2.968* (1.261)	-3.491** (1.230)	-3.762** (1.170)
neighbourhood solid fuel use			-9.188*** (2.739)	-6.127* (2.628)	-6.424* (2.612)	-6.088* (2.450)
female	-0.650 (0.936)	0.543 (0.828)	0.584 (0.828)	-0.367 (0.877)	-0.576 (0.939)	-0.859 (0.908)
rural	-2.691* (1.131)	-0.533 (1.367)	1.301 (1.898)	1.320 (1.816)	1.384 (1.804)	1.382 (1.672)
age	-0.124*** (0.021)	-0.134*** (0.021)	-0.134*** (0.021)	-0.124*** (0.021)	-0.127*** (0.024)	
assets					-0.070 (0.086)	
education					-0.048 (0.092)	
state fixed effects	X	X	X	X	X	X
age in years (as dummies)						X
asset count (as dummies)						X
years of education (as dummies)						X
constant	74.108*** (2.044)	76.179*** (1.993)	77.875*** (2.125)	78.248*** (2.140)	80.334*** (2.556)	74.768*** (2.619)
r-squared	0.048	0.053	0.053	0.060	0.060	0.090
Observations	9551	9551	9551	9551	9551	9551

Source: WHO SAGE 2007-08

Note: Robust standard errors, clustered at the level of the primary sampling unit, in parentheses. + p<.1, * p<.05, ** p<.01, *** p<.001. Outcome: FEV1% (FEV1s/FVC as a percentage) is the ratio of the volume of air one can exhale in 1 second to the total volume of air one can exhale after inhaling as deeply as possible.

Table A2: Robustness: Additional controls for household and neighbourhood socioeconomic status

	(1)	(2)	(3)	(4)
smoke tobacco	-3.764*** (1.04)	-3.792*** (1.04)	-3.774*** (1.04)	-3.770*** (1.04)
household uses solid fuels	-3.762** (1.17)	-3.577** (1.15)	-3.352** (1.14)	-3.363** (1.14)
neighbourhood solid fuel use	-6.088* (2.45)	-6.397* (2.59)	-7.936** (2.78)	-7.492* (3.18)
log monthly expenditure		1.012+ (0.60)	0.838 (0.61)	0.839 (0.61)
log monthly expenditure squared		-0.667 (1.21)	0.255 (1.32)	0.284 (1.31)
average neighbourhood assets			-0.342 (0.24)	-0.385 (0.27)
average neighbourhood education				0.114 (0.38)
demographic controls	X	X	X	X
place controls	X	X	X	X
ses controls	X	X	X	X
constant	74.768*** (2.62)	72.409*** (10.95)	71.927*** (10.83)	71.393*** (10.80)
observations	9551	9551	9551	9551

Source: WHO SAGE 2007-08

Note: Robust standard errors, clustered at the level of the primary sampling unit, in parentheses. + $p < .1$, * $p < .05$, ** $p < .01$, *** $p < .001$. Outcome: FEV1% (FEV1s/FVC as a percentage) is the ratio of the volume of air one can exhale in 1 second to the total volume of air one can exhale after inhaling as deeply as possible.

Table A3: Placebo tests for smoking, household solid fuel use, and neighbourhood solid fuel use

	A. Smoking tobacco		B. Household solid fuel use		C. Neighbourhood solid fuel use			
	(1)	(2)	(1)	(2)	(1)	(2)		
chew tobacco	-0.28 (0.86)		hh owns toilet	0.742 (0.94)	neighbourhood open defecation	-0.178 (1.95)		
drink alcohol		-1.318 (1.34)	hh electrified		neighbourhood electrification	-0.83 (2.05)		
smoke tobacco	-3.772*** (1.10)	-3.522*** (1.06)	hh uses solid fuels	-5.501*** (1.05)	-5.766*** (1.07)	neighbourhood solid fuel use	-7.309*** (2.20)	-7.852*** (1.83)
demographic controls	X	X	demographic controls	X	X	demographic controls	X	X
place controls	X	X	place controls	X	X	place controls	X	X
ses controls	X	X	ses controls	X	X	ses controls	X	X
constant	77.408*** (2.99)	77.416*** (2.95)	constant	82.242*** (3.16)	82.429*** (3.32)	constant	83.252*** (3.34)	84.130*** (4.00)
observations	9551	9551	observations	9551	9551	observations	9551	9551

Source: WHO SAGE 2007-08

Note: Robust standard errors, clustered at the level of the primary sampling unit, in parentheses. + p<.1, * p<.05, ** p<.01, *** p<.001. All regressions are weighed using national individual weights. Outcome: FEV1% (FEV1s/FVC as a percentage) is the ratio of the volume of air one can exhale in 1 second to the total volume of air one can exhale after inhaling as deeply as possible. hh: household.

Table A4: Falsification tests and tests with alternative measures of respiratory health: Association between other health outcomes and exposures

test: specification:	falsification		alternative measures of respiratory health		
	ols		logit		
	FVC	height	FEV1s	have COPD (FEV1% < 70%)	self-reported lung disease
outcome:	(1)	(2)	(3)	(4)	(5)
smoke tobacco	-0.038 (0.058)	0.002 (0.004)	-0.189*** (0.030)	1.422* (0.195)	1.526* (0.282)
household uses solid fuels	-0.050 (0.059)	-0.000 (0.004)	-0.139*** (0.037)	1.467* (0.220)	1.919* (0.505)
neighbourhood solid fuel use	-0.008 (0.075)	0.002 (0.005)	-0.162** (0.061)	1.502* (0.290)	1.966* (0.668)
demographic controls	X	X	X	X	X
place controls	X	X	X	X	X
ses controls	X	X	X	X	X
height			X		
constant	2.765*** (0.414)	1.656*** (0.014)	0.755 (0.565)		
observations	9551	9551	9551	9551	9551

Source: WHO SAGE 2007-08

Note: Robust standard errors, clustered at the level of the primary sampling unit, in parentheses. + p<.1, * p<.05, ** p<.01, *** p<.001. Outcomes: FVC, or Forced Vital Capacity is the total amount (measured in liters) of air breathed out in the spirometry test. Height is measured in meters. FEV1s or Forced Expiratory Volume in 1 second (measured in liters) is the total amount of air breathed out in the first second of the spirometry test. Respondents are classified as having COPD if FEV1% was below 70%, based on GOLD criteria. Respondents classified as having lung disease based on self-reported symptoms according to WHO SAGE criteria, of having shortness of breath or experiencing coughing or wheezing for more than ten minutes and coughing sputum or phlegm for most days of a month for the past three months in a year.